



ESTIMATION OF EXPOSURE LEVELS OF TERRESTRIAL BIOTA AND RADIATION EXPOSURE AROUND IPEN'S FACILITIES

Cavalcante^a F., Nisti^a M.B., Saueia^a C.H.R.

^a Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN – SP)

Postal code 05508-000, São Paulo, SP, Brazil

mbnisti@ipen.br

ABSTRACT

Humans are exposed to natural radiation; soil is a major source of external and internal exposure of radiation. The aim of this study is to evaluate the exposure levels of terrestrial biota and to estimate the radiation exposure around Instituto de Pesquisas Energéticas e Nucleares (IPEN) facilities. The ERICA Tool was used to calculate the exposure levels of terrestrial biota; the estimation of radiation exposure for humans was determined using a model proposed by UNSCEAR and Excess Lifetime Cancer Risk. Six soil points were collected and their activity concentrations were measured by gamma spectrometry. Two soil points showed a risk coefficient greater than 1, suggesting that the screening dose ratio of $10\mu\text{Gy h}^{-1}$ might be exceeding for the most exposed species, namely lichen and bryophytes, even though the activity concentration values of the analyzed radionuclides showed no evidence of soil contamination due to the atmospheric discharges of the IPEN facilities. Thus, the radioactive discharges in the soil from all facilities are negligible. Hence, the authors concluded that the ERICA Tool can be useful in assisting environmental radiological monitoring program for decision-making, especially regarding: points collected, sample types and sampling frequency.

Keywords: estimation of radiation exposure; ERICA Tool; biota; gamma spectrometry.

1. INTRODUCTION

Biota and human are exposed to natural radiation from many different sources. The external exposure from the soil is associated with gamma radiation and internal exposure with radon inhalation, soil is a major source of exposure of radiation, and the exposures of radiation are different in each region [1].

In the past, the levels of exposure to ionizing radiation was mainly focused on humans, considering that biota and the environment were also protected if human beings were adequately protected. In the last decades, this statement was proven to fail and is no longer accepted [2]. Exposure and radiological risk to biota from different ecosystems can be assessed using different risk models, such as the RESRAD-Biota and the ERICA Tool.

The Erica Integrated Approach [3] was developed by the European Union to assess the effects of radionuclides in the environment and to support decision making. The software operates in three different Tiers and provides estimation on absorbed doses (internal and external) to reference organisms from different ecosystems and perform risk characterization based on activity concentration in the environment and in biota whole body.

The aim of this study is to evaluate the exposure levels of terrestrial biota and to estimate the radiation exposure around Instituto of Pesquisas Energéticas e Nucleares (IPEN) facilities. The ERICA Tool was used to calculate the exposure levels of terrestrial biota; the estimation of radiation exposure for humans was determined using theoretical models proposed by UNSCEAR and Excess Lifetime Cancer Risk. Soil samples were collected and their activity concentrations were measured by gamma spectrometry.

Estimation of Radiation Doses (Absorbed Dose Rate in air, Annual Effective Dose) and Excess Lifetime Cancer Risk (ELCR) were calculated using theoretical models [1, 4].

2. MATERIALS AND METHODS

The IPEN is located in the city of Sao Paulo – Brazil and comprises several nuclear and radioactive facilities, including a research reactor, cyclotrons and a radioisotope and radiopharmaceutical production plant. Gaseous and Liquid radioactive effluents are acutely monitored before released into the environment and verified by the Environmental Radiological Monitoring Program (PMRA).

2.1. Data on soil activity concentrations from natural radionuclides

Nisti et al. [5] determined the activity concentrations from natural radionuclides in soil from different locations in IPEN, giving evidence of no soil contamination due to the atmospheric releases from IPEN facilities. Soil samples were measured by gamma spectrometry with a hyper-pure germanium detector Canberra model GX2518, 25% relative efficiency, effective resolution of 1.8 keV on the 1332 keV ^{60}Co with associated electronics. Table 1 presents the results obtained in the referred paper:

Table 1: Average concentrations of ^{226}Ra , ^{210}Pb , ^{232}Th , ^{228}Th and ^{40}K in soil samples (Bq kg^{-1}) and sampling location from Nisti et al. (2015)

Sampling coordinates	Concentration (Bq kg^{-1})				
	^{226}Ra	^{210}Pb	^{232}Th	^{228}Th	^{40}K
1 23°33'56.66"S-46°44'07.04"O	43 ± 3	61 ± 12	92 ± 4	101 ± 4	179 ± 8
2 23°33'55.64"S-46°44'05.63"O	52 ± 1	47 ± 5	124 ± 5	134 ± 7	94 ± 10
3 23°33'59.69"S-46°44'15.48"O	42 ± 1	43 ± 6	83 ± 2	90 ± 5	200 ± 11
4 23°33'46.15"S-46°44'13.36"O	40 ± 2	62 ± 10	83 ± 7	90 ± 1	143 ± 13
5 23°33'48.21"S-46°44'16.35"O	39 ± 2	51 ± 3	70 ± 2	79 ± 6	204 ± 11
6 23°33'41.26"S-46°44'28.92"O	54 ± 5	59 ± 6	116 ± 2	127 ± 6	185 ± 18
UNSCEAR [1]	17-60		11-64		140-850
Peres [8]	1-61.8	<20-121	8-82	4.8-120	15.3-516

2.2. Risk characterization for terrestrial biota using Erica Tool

Data on radionuclide concentrations on soil were used as input to calculate the Risk Quotient (RQ) for all terrestrial reference organisms. The assessment was run using Tier 1, once this is more conservative and only requires media concentration activities.

Whenever the calculations of RQ's present a value equal to or higher than 1, it indicates that there is a significant probability that the activity concentration of a particular radionuclide exceeds the screening dose value (10 μ Gy/h) for the most exposed organism. The Tool suggests that the user carry on with the assessment, using Tier 2 or Tier 3.

The assessment using Tier 2 requires the activity concentrations in biota whole body in order to estimate the total absorbed dose (from internal and external sources). Once estimated, the Tool employs these results to calculate a new value for the Risk Quotient.

According to Brown J.E. et al (2008), if adequate measured values of activity concentrations in biota whole body are not available, one can infer them using the Concentration Ratio (CR) given by the following equation:

$$CR = \frac{\text{Activity concentration in biota whole body (Bq.kg}^{-1} \text{ fresh weight)}}{\text{Activity concentration in soil (Bq.kg}^{-1} \text{ dry weight)}} \quad (1)$$

There is a wide set of CR's values stored in the Tool for each radionuclide and reference organism.

The reference organisms for terrestrial biota evaluated in this paper were: bird, flying insects, grasses and herbs, lichen and bryophytes, shrub and tree.

2.3. Estimation of Radiation Doses and Excess Lifetime Cancer Risk

The ^{226}Ra , ^{232}Th and ^{40}K activity concentrations of soil samples were used for the calculation of outdoor external absorbed dose rate in air at 1m above the ground surface to the population [1]. The absorbed dose rate in air was obtained by the equation (2):

$$DR = [(CRa \times CFRa) + (CTh \times CFTTh) + (CK \times CFK)] \quad (2)$$

Where:

DR is external absorbed dose rate (nGy h^{-1}),

CRa is the activity concentrations of ^{226}Ra (Bq kg^{-1}),

CTh is the activity concentrations of ^{232}Th (Bq kg^{-1}),

Ck is the activity concentrations of ^{40}K (Bq kg^{-1}),

CFRa is conversion factor of ^{226}Ra 0.462 (nGy h^{-1} per Bq kg^{-1}),

CFTTh is conversion factor of ^{232}Th 0.604 (nGy h^{-1} per Bq kg^{-1}),

CFk is conversion factor of ^{40}K 0.0417 (nGy h^{-1} per Bq kg^{-1}),

The conversion factors were defined by the UNSCEAR [6].

The annual effective dose (outdoor) to the population was calculated using the following equation 3:

$$AED = (DR \times T \times OF \times CF) \quad (3)$$

Where:

AED is annual effective dose (outdoor) (mSv y^{-1}),

DR is external absorbed dose rate (nGy h^{-1}),

T is annual average time for exposure to radiation (h),

OF is outdoor occupancy factor 0.2,

CF is conversion factor 0.7 (Sv Gy^{-1}).

The occupancy factor and conversion factor was proposed by the UNSCEAR [6].

The Excess Lifetime Cancer Risk is calculated using the following equation 4.

$$ELCR = AED \times DL \times RF \quad (4)$$

Where:

ELCR is Excess Lifetime Cancer Risk,

AED is annual effective dose (outdoor) (mSv y⁻¹),

DL is duration of life 70 (years),

RF is risk factor 0.05x10⁻³ (Sv⁻¹).

The duration of life and risk factor for stochastic effect was defined by the ICRP [7].

3. RESULTS AND DISCUSSION

3.1. Risk analysis and absorbed dose rates for terrestrial biota

Tier 1 was initially used to calculate the RQ for terrestrial organisms, using data from Table 1 as input. The results for each studied radionuclide and the most exposed organism are presented in Table 2, using the highest value for each radionuclide concentration.

Table 2: Risk Quotient for each radionuclide and limiting reference organism

Isotopes	RQ [unitless] (Mn – max)	Limiting Reference Organism
Ra-226	1.88 E0	Lichen & Bryophytes
Pb-210	9.92 E-3	Lichen & Bryophytes
Th-232	3.99 E-1	Lichen & Bryophytes
Th-228	3.47 E0	Lichen & Bryophytes

Calculations have shown that the concentrations of Ra-226 and Th-228 measured in soil may provide absorbed doses higher than the screening dose value (10µGy/h) for the most exposed organism (lichen and bryophytes). The Tool indicates the need to further investigate these concentrations, which was done using Tier 2.


As mentioned before, Tier 2 requires the activity concentrations in biota whole body in order to estimate the total absorbed dose (from internal and external sources) and calculates a new set of Risk Quotient values. Activity concentration in biota whole body were not available in this work, therefore, the authors addressed to the default values of CR within the Tool.

Using equation 1 and default values of CR, the activity concentration in biota whole body was than inferred for selected organisms (Table 3) and are shown next in Figure 1:

Table 3: Whole body activity concentration for terrestrial biota, inferred from default CR values and measured soil activity concentrations

Whole Body Activity Concentrations (Bq kg ⁻¹ f.w.)						
Isotope	Lichen & Bryophytes	Grasses & Herbs	Bird	Flying insects	Shrub	Tree
Ra-226	38.34	9.4	1.9	2.3	17.0	0.6
Pb-210	160.0	7.4	3.8	25.0	20.0	4.31
Th-228	51.0	21.0	0.052	0.68	8.2	0.17
Th-232	47.0	2.0	0.048	0.63	7.6	0.16

Figure 1. Activity concentration in organism whole body, due to internal and external exposures of radionuclides in soil.



In agreement with the results obtained in Tier 1, the activity concentration in the most exposed organism, lichen and bryophytes, are higher, specially for Pb-210, as expected. Once Rn-222 is easily released to the air from soil surfaces, some of their radioactive daughters (such as Po-210) can be deposited on the surface of vegetation.

The estimated values presented in Table 3 were used as input to calculate the internal (Table 4) and external (Table 5) dose rates and a new set of RQ's (Table 6) for each radionuclide and exposed organism.

Table 4: Internal absorbed doses for selected organisms

Internal absorbed doses ($\mu\text{Gy h}^{-1}$)						
Isotope	Lichen & Bryophytes	Grasses & Herbs	Bird	Flying insects	Shrub	Tree
Ra-226	5.31	1.28	0.27	0.31	2.32	0.084
Pb-210	0.031	0.002	9.88E-4	0.005	0.0045	0.0011
Th-228	9.451	3.881	0.01	0.126	1.51	0.03
Th-232	1.081	0.04	0.001	0.014	0.17	0.004

Table 5: External absorbed doses for selected organisms

External absorbed doses ($\mu\text{Gy h}^{-1}$)						
Isotope	Lichen & Bryophytes	Grasses & Herbs	Bird	Flying insects	Shrub	Tree
Ra-226	0.018	0.018	0.012	0.0189	0.01728	0.014
Pb-210	1.80E-5	2.48E-5	1.736E-5	1.80E-5	1.24E-5	8.06E-6
Th-228	0.039	0.04	0.03886	0.03886	0.036	0.03082
Th-232	5.42E-6	1.36E-5	5.33E-6	5.45E-6	6.2E-6	2.604E-6

Table 6: Expected RQ's for each selected organism.

Organism	Total Dose Rate per organism ($\mu\text{Gy h}^{-1}$)	Screening Value ($\mu\text{Gy h}^{-1}$)	Risk Quotient (expected) (unitless)
Lichen & Bryophytes	1.59E1	1.00E1	1.59E0
Grasses & Herbs	5.27E0	1.00E1	5.27E-1
Bird	3.41E-1	1.00E1	3.41E-2
Flying insects	5.17E-1	1.00E1	5.17E-2
Shrub	4.07E0	1.00E1	4.07E-1
Tree	1.67E-1	1.00E1	1.67E-2

The Tool points out three different results: for lichen & bryophytes, the screening dose rate is exceeded and requires further investigation; for grasses & herbs and shrubs, there is a significant probability that the screening dose rate might be exceeded; for birds, flying insects and trees, the probability that the screening dose rate is exceeded is low. Further investigation (including area characterization and environmental sampling) in the area with the highest soil activity concentration is currently under consideration and will be addressed in future papers.

3.2. Radiation Doses and Excess Lifetime Cancer Risk

Two scenarios were used to estimate the assumption of annual average time for exposure to radiation of Annual Effective Dose of the population in IPEN. The first scenario, more conservative, considered the time of 8,766 hours (Table 7). The second scenario, more realistic, considered the hours that the individual (worker) is inside IPEN, estimated at 2,277 hours (Table 8).

Table 7: Estimation of Radiation Doses (Absorbed Dose Rate in air (DR), Annual Effective Dose (AED)) and Excess Lifetime Cancer Risk (ELCR) (first scenario).

Sampling coordinates	DR (nGy h ⁻¹)	AED (mSv y ⁻¹)	ELCR
1 23°33'56.66"S-46°44'07.04"O	83	0.10	0.36x10 ⁻³
2 23°33'55.64"S-46°44'05.63"O	103	0.13	0.44 x10 ⁻³
3 23°33'59.69"S-46°44'15.48"O	78	0.10	0.33 x10 ⁻³
4 23°33'46.15"S-46°44'13.36"O	75	0.09	0.32 x10 ⁻³
5 23°33'48.21"S-46°44'16.35"O	69	0.08	0.30 x10 ⁻³
6 23°33'41.26"S-46°44'28.92"O	103	0.13	0.44 x10 ⁻³
mean ± st. dev.	85±15	0.10±0.02	(0.36±0.06) x10 ⁻³

The results obtained for the absorbed dose in air from soil varied from 69 to 103 nGy h⁻¹ in soil samples of IPEN, with an average value of 85 ± 15 nGy h⁻¹.

The Annual Effective Dose (outdoor) from soil varied from 0.08 to 0.13 mSv y⁻¹, with an average value of 0.10 ± 0.02 mSv y⁻¹.

The Excess Lifetime Cancer Risk (outdoor) from soil varied from 0.30x10⁻³ to 0.44x10⁻³, with an average value of (0.36 ± 0.06)x10⁻³.

Table 8: Estimation of Radiation Doses (Absorbed Dose Rate in air (DR), Annual Effective Dose (AED)) and Excess Lifetime Cancer Risk (ELCR) (second scenario).

Sampling coordinates	DR (nGy h ⁻¹)	AED (mSv y ⁻¹)	ELCR
1 23°33'56.66"S-46°44'07.04"O	83	0.03	0.09x10 ⁻³
2 23°33'55.64"S-46°44'05.63"O	103	0.03	0.11x10 ⁻³
3 23°33'59.69"S-46°44'15.48"O	78	0.02	0.09x10 ⁻³
4 23°33'46.15"S-46°44'13.36"O	75	0.02	0.08x10 ⁻³
5 23°33'48.21"S-46°44'16.35"O	69	0.02	0.08x10 ⁻³
6 23°33'41.26"S-46°44'28.92"O	103	0.03	0.11x10 ⁻³
mean ± st. dev.	85±15	0.03±0.01	(0.01±0.02) x10 ⁻³

The Annual Effective Dose (outdoor) from soil varied from 0.02 to 0.03 mSv y⁻¹, with an average value of 0.03 ± 0.01 mSv y⁻¹.

The Excess Lifetime Cancer Risk (outdoor) from soil varied from 0.08x10⁻³ to 0.11x10⁻³, with an average value of (0.10 ± 0.02) x 10⁻³.

The results obtained for the absorbed dose in air (outdoor) are in good agreement with the value reported from UNSCEAR [1] for the range worldwide of 18 to 93 nGy h⁻¹.

The Annual Effective Dose (External terrestrial radiation - Outdoors) and Excess Lifetime Cancer Risk in the present studies are of the same order of magnitude of the mean worldwide of 0.07 mSv y⁻¹ and 0.29x10⁻³, respectively [1, 4, 6].

4. CONCLUSION

Risk Quotients calculations using Tier 1 have shown that the concentrations of Ra-226 and Th-228 measured in soil may provide absorbed doses higher than the screening dose value ($10\mu\text{Gy h}^{-1}$) for the most exposed organism (lichen and bryophytes). Using Tier 2, the Tool provided a new set of Risk Quotients as well as the internal and external dose rates for selected organisms. For lichen & bryophytes, the screening dose rate is exceeded and requires further investigation; for grasses & herbs and shrubs, there is a significant probability that the screening dose rate might be exceeded.

According to these results, the authors believe that the Tool can be used to justify further investigation of the area with the highest soil activity concentration. Therefore, the authors conclude the ERICA Tool can be useful in assisting environmental radiological monitoring program for decision-making, especially regarding: points collected, sample types and sampling frequency.

The results obtained for Estimation of Radiation Doses (Absorbed Dose Rate in air (DR), Annual Effective Dose (AED)) and Excess Lifetime Cancer Risk (ELCR) in the soil samples for both scenarios; indicate that the exposure around IPEN facilities is of the same order of magnitude of the mean worldwide.

Finally, the results of this paper can be used for a database on soil radioactivity in the São Paulo city.

REFERENCES

1. UNSCEAR- United Nations Scientific Committee on the Effects of Atomic Radiation. Sources, Effects and Risks of Ionization Radiation; Report to the General Assembly, with Annexes, New York, 2000.
2. Vives i Batlle, J., Balonov, M., Beaugelin-Seiller, K., Beresford, N., Brown, J., Cheng, J. J., Coplestone, D., Doi, M., Filistovic, V., Golikov, V., et al. (2007). Inter-comparison of absorbed dose rates for non-human biota. *Radiation and Environmental Biophysics*, 46 (4): 349-373. doi: 10.1007/s00411-007-0124-1.
3. Brown, J. E., Alfonso, B., Avila, R., Beresford, N. A., Coplestone, D., Prohl, G. & Ulanovsky, A. (2008). The ERICA Tool. *J Environ Radioact*, 99 (9): 1371-83. doi: 10.1016/j.jenvrad.2008.01.008.
4. Taskin, H., Karavus, M., Ay, P., Topuzoglu, A., Hidiroglu, S., Karahan, G. Radionuclide concentrations in soil and lifetime cancer risk due to the gamma radioactivity in Kirklareli, Turkey. *Journal of Environmental Radioactivity*, v. 100, No. 1, p. 49-53. 2009.
5. Nisti M. B., SAUEIA, C. H. R., MAZZILLI, B. P. Chapter 20. Determination of Radioactivity Levels and ^{222}Rn Exhalation Rates in Soil Around Ipen Facilities. Special Publications. 1ed.: Royal Society of Chemistry, v. 1, p. 159-163, 2015.
6. UNSCEAR- United Nations Scientific Committee on the Effects of Atomic Radiation “Effects of ionizing radiation”: report to the General Assembly, with scientific annexes. Vol. 1. United Nations Publications. 2008.
7. ICRP-International Commission on Radiological Protection. The 2007 Recommendations of the International Commission on Radiological Protection. v. 37 n. 2-4. ICRP Publication 103. 2007.
8. PERES, A.C. Modelo para o estabelecimento de valores orientadores para elementos radioativos no solo. 2007. Tese (Doutorado em Tecnologia Nuclear) – Instituto de Pesquisas Energéticas e Nucleares – IPEN/CNEN/SP, São Paulo.