



Iodine determination in edible algae species using epithermal neutron activation analysis methodology

Albuquerque^a C.R., Maihara^a V.A., Silva^a P.S.C.

^a Instituto de Pesquisas Energéticas e Nucleares, Centro do Reator de Pesquisas /Laboratório de Análise por Ativação
Neutrônica, 05508-000, São Paulo, SP, Brazil

calbuuquerque@gmail.com

ABSTRACT

Iodine is one of the essential trace elements of much interest in nutritional research being responsible for the production of the thyroid hormones, which has great importance for human metabolism. Edible algae species accumulate iodine from seawater and are considered a good dietary source of this nutrient. The Epithermal Neutron Activation Analysis (ENAA) was applied to determine the iodine concentration in edible algae species and derivatives. Twenty-one samples for 4 edible algae species *Porphyra umbilicalis* (common name: Nori), *Hijikia fusiforme* (Hijiki), *Undaria pinnatifida* (Wakame) and *Laminaria* spp. (Kombu) and 3 samples of the derivative sample, ágar, were analyzed. The results pointed out a great variability of I concentration in these species. The highest I concentration obtained was 9324 ± 113 mg/kg in *Laminaria* spp. (Kombu). *Hijikia fusiforme* (Hijiki) also presented high values (1803 ± 86 mg/kg). The lowest I content was obtained in agar samples (3.18 ± 0.29 mg/kg). All analyzed algae samples exceed the recommended value (150 µg/day) and an ingestion of 1 g/day of *Laminaria* spp. (Kombu) exceed the maximum tolerable limit (1100 µg/day) and its consumption must be controlled to avoid possible health problems.

Keywords: algae , iodine, epithermal neutron activation analysis.

1. INTRODUCTION

Iodine is a trace element essential for the body because it is the main constituent of thyroid hormones. The thyroid gland is responsible for cell growth, brain development, and maintaining body temperature. About 99% of iodine available in the body is located in the thyroid gland [1,2].

Goitre is one of the most important disorders caused due to iodine deficiency, apart from hypothyroidism and cretinism. To overcome iodine deficiency in the diet of the population, mainly in underdeveloped countries, iodine supplementation in the diet has been performed, for example iodination on the cooking salt. On the other hand, iodine supplementation programs should be monitored to avoid the presence of diseases caused by its excess such as hyperthyroidism [3].

In food, iodine concentrations and total iodine dietary intake differ from one region to another, due to geochemical characteristics, soil, and cultural conditions that interfere with food choice [1].

Algae are excellent sources of minerals and may present higher values in relation to vascular plants and terrestrial animals [4] since they absorb inorganic substances from the environment and store them [5,6].

Algae are also used as bioindicators of environmental contamination due to their morphological and physiological characteristics, as they have a high capacity to absorb and store certain potentially toxic elements [7,8] The group of brown algae, such as *Hijikia fusiforme* (Hijiki), *Undaria pinnatifida* (Wakame) and *Laminaria* spp. (Kombu) has greater capacity to store iodine in relation to the group of red algae (*Porphyra umbilicalis* (Nori) [7].

The species red algae are the most commercially used for food and feed, for the extraction of Agar (manufacture of gums, laxatives, culture medium for bacteria) and for the extraction of carrageenan (use in dairy products, in the manufacture of jellies and jellies, as a thickener in soups, sauces and as a non-caloric substitute for starch and fat)[9].

In many countries, food industries employ a wide range of algae derivatives, which contains high levels of fiber, fatty acids, polyunsaturated polysaccharides, minerals, vitamins and antioxidants. The algae and its derivatives also have economic impact in several sectors, such as aquaculture, pharmaceutical industry, biomedicine, veterinary medicine, cosmetics industry and public health [8,10]. Studies about iodine concentration in algae are increasing due to question

related to public health, since there are several pathologies caused by the deficiency and excess consumption of this element.

The aim of this study was to determine the concentration of iodine in brown and red algae and its derivative agar, and check if the found levels are in agreement with the recommend limits for human health.

2. MATERIALS AND METHODS

2.1. Sample Preparation

Twenty-four samples from four species of commercially edible algae available to consumers were purchased from supermarkets around São Paulo city. All edible algae samples were imported from 4 different countries. The samples were ground and homogenized using a household mixer, fitted with titanium blade, until they acquire a powder consistency. Table 1 shows the different species of algae (with its commercial name), its origins and the number of each species acquired.

Table 1: Species of algae and derivates, countries of origin and number of samples

Algae and Derivates	Country	Number of samples
<i>Porphyra umbilicalis</i> (Nori)	China	1
<i>Porphyra umbilicalis</i> (Nori)	Korea	2
<i>Porphyra umbilicalis</i> (Nori)	Japan	2
<i>Porphyra umbilicalis</i> (Nori)	USA	1
<i>Hijikia fusiforme</i> (Hijiki)	Japan	2
<i>Hijikia fusiforme</i> (Hijiki)	China	2
<i>Hijikia fusiforme</i> (Hijiki)	Korea	1
<i>Undaria pinnatifida</i> (Wakame)	China	3
<i>Undaria pinnatifida</i> (Wakame)	Korea	2
<i>Laminaria sp</i> (Kombu)	Korea	1
<i>Laminaria sp</i> (Kombu)	Japan	2
<i>Laminaria sp</i> (Kombu)	China	2
Agar	Brazil	3

2.2. Epithermal Neutron Activation Analysis (ENAA)

Iodine determination was carried out using Epithermal Neutron Activation Analysis (ENAA) [11]. The principle of ENAA consists in irradiating sample wrapped with a cadmium sheet to shield out the thermal neutrons and only epithermal and rapid neutrons target the sample. Some radioisotopes have high capture resonance integral, like ^{128}I , and they are greatly enhanced by epithermal activation. Thereby it is possible reduce possible interferences due to activation of elements with high cross section for thermal neutrons, such as ^{24}Na , ^{38}Cl present in the algae samples.

The following sample and standard preparation and the irradiation conditions are the in-house procedure [12]. About 50 mg of the algae samples and reference materials were weighed in acid cleaned polyethylene bags and iodine standards were prepared by pipetting the KIO_3 solution (6.8 μg of I) on Whatman 40 paper and dried at room temperature.

The algae samples, reference materials and Iodine standards were wrapped with a cadmium sheet with thickness of 2 mm and then irradiated for 30 seconds under a neutron flux of $10^{12} \text{ n cm}^{-2} \text{ s}^{-1}$ in the IEA-R1 pneumatic system of IPEN - CNEN/SP. After decay time of about 600 seconds, the radioisotope ^{128}I was determined by measuring its gamma ray transition of 442.9 keV by gamma spectrometry.

Gamma ray measurements were performed using a GC2018 Canberra HPGe detector coupled to a Canberra DSA-1000 multichannel analyzer. The spectra were collected and processed using a Canberra Genie 2000 version 3.1 spectroscopy software.

2.3 Quality control

To validate the ENAA method in respect of its precision and accuracy two standard reference materials (SRM) from National Institute of Standards and Technology (NIST): Whole milk (NIST RM 8435) and Whole egg powder (NIST RM 8415) were analyzed. The determined concentrations

were compared with the value of their certificates by means of the E_n -score method¹ [13] Figure 1 shows the E_n -score values for the element iodine in the SRM.


The Table 2 shows relative errors less than 9% and relative standard deviations below 12%, indicating good precision and accuracy of the data.

Table 2 - Iodine concentration results obtained for Whole Milk Powder (RM 8435) and Whole Egg Powder (RM 8415) reference materials.

CRM	Determined concentration(M \pm DP) ^a mg/kg	SD Rel%	ER%	Certificate concentration (mg/kg)
WHOLE MILK POWDER	2.50 \pm 0.16	6.4	8.7	2.3 \pm 0.4
WHOLE EGG POWDER	2.08 \pm 0.24	11.5	5.6	1.97 \pm 0.50

The E_n -score values obtained for iodine in the reference materials showed in Figure 1 are less than 1, so they are accurate with the values of the certificates.

Figure 1: E_n -score values for Iodine determined in the RM



¹ The E_n -score test represents a normalized Gaussian centered at 0, and values between -2 and 2 for this test indicate that within a 95% confidence interval the value is satisfactory. ^a arithmetic mean and the standard deviation of 3 determinations.

3. RESULTS AND DISCUSSION

3.1 Results of edible seaweeds

Table 3 presents the average results for I in algae samples, obtained in duplicates. All the results presented are related to the dry weight (dw) of the samples.

The results showed that there is a great variability between species in relation to the iodine concentration. The values of relative standard deviation (RSD) ranged from 0.9 to 17%. The higher RSD observed in all samples of *Undaria* sp (Wakame) is probably related to the high Na concentrations (from up to 7 to 8.8%) and that the R_{Cd} (cadmium ratio) for sodium is 58 a significant amount of this element must be activated. According to the literature the brown algae present higher I level than the others due its greater affinity for this element. In this study the same was observed as shown in the Figure 2.

In Nagataki study in 2008 [14], the algae of the species *Laminaria* spp (Kombu) presented contents of 0.3% of I in dry weight (dw) and *Undaria* sp (Wakame) presented levels of 0.02-0.03% (dw), for the red algae *Porphyra* (Nori) the result was <0.01% (dw). Fukushima and Chatt, 2012 [15], also obtained high concentrations of I in brown algae of the species *Undaria pinnatifida* (Wakame), and *Hijikia fulsiforme* (Hijiki). Hijiki presented the I content of 514 ± 6 mg/kg, and Wakame 156 ± 3 mg/kg. In relation to other species of algae these are in fact those with higher I content.

In this study the highest determined concentrations for the brown algae were: 9324 ± 113 mg/kg for *Laminaria* spp (Kombu), 1803 ± 86 mg/kg for *Hijikia fulsiforme* (Hijiki) and 430 ± 74 mg/kg for the *Undaria pinnatifida* (Wakame) can be observed in the Figure 2. For the red algae *Porphyra umbilicalis* the maximum concentration was 103 ± 1 mg/kg (Figure 3). The concentrations for *Ágar*, a derivative of red algae, presented low levels of I, ranging from 3.18 ± 0.29 mg / kg up to 7.6 ± 0.5 mg / kg as can be shown on Figure 4.

It is important to notice that the daily intake of iodine according to DRI (dietary recommended intake) for adults is up to 100 to 150 μ g/day, and the maximum tolerable limit (UL) is 1100 μ g/day [15]. The determined results of iodine concentration in brown algae varied from 116 to 9324 mg/kg

which means that the ingestion of 1 g of brown algae (in case of Kombu) can exceed 62 times the DRI and 8 times the UL (for the higher I concentration).

Table 3 - Results of I concentration in seaweed samples by ENAA in dry basis

Specie	Origen	I mg/kg Median \pm SD ^a	RSD(%)
<i>Porphyra umbilicalis</i> (Nori)	China	44.1 \pm 8.9	20.1
	Korea	33.4 \pm 4.0	12.1
	Korea	33.4 \pm 2.4	7.2
	Japan	114.09 \pm 0.02	0.01
	USA	54.5 \pm 9.7	17.7
<i>Hijikia fusiforme</i> (Hijiki)	China	1975 \pm 94	4.8
	China	1494 \pm 252	16.9
	Korea	1365 \pm 13	0.9
	Japan	734 \pm 12	1.6
	Japan	836.8 \pm 8.9	1.1
<i>Laminaria sp</i> (Kombu)	China	2826 \pm 301	10.8
	China	259 \pm 29	11.1
	Korea	9962 \pm 121	1.2
	Japan	6810 \pm 435	6.4
	Japan	4761 \pm 195	4.1
<i>Undaria pinnatifida</i> (Wakame)	China	126 \pm 12	9.7
	China	468 \pm 80	17.2
	China	267 \pm 45	16.8
	Korea	398 \pm 66	16,6
	Korea	566 \pm 109	19.3
Ágar I	Brazil	8.29 \pm 0.08	0.9
Ágar II	Brazil	10.6 \pm 2.3	21
Ágar III	Brazil	4.03 \pm 0.73	11.1

Figure 2: Values of the Iodine concentration in brown algae.




Figure 3: Values of the Iodine concentration in *Porphyra umbilicalis* (Nori).





Figure 4: Values of the Iodine concentration in Ágar.

4. CONCLUSION

The ENAA technique proved adequate for the determination of iodine in seaweed samples. The results showed that edible marine algae can be a great source of iodine, especially brown algae, and can be used in food to meet some nutritional needs, but consumption must be controlled to avoid possible health problems. In relation to the agar a derivative of the red algae, it was possible to conclude that it does not present risks for the feeding, in relation to the iodine element.

ACKNOWLEDGMENT

The authors would like to thank the Institute of Energy and Nuclear Research (IPEN-CNEN/SP) and the CNPq for the scholarship granted (Process number 133916/2017-0).

REFERENCES

- [1] JACOB, M.; BRITO, N. Suplementação de iodo na gravidez: qual a importância? **Rev. Port. saúde pública**, v. 3, n. 1, p. 107–119, 2015.
- [2] HALDIMANN, M., ALT, A., BLANC, A., BLONDEAU, K. Iodine content of food groups. **J Food Compos Anal**, v. 18, n. 6, p. 461-471, 2005.
- [3] G. S. HENRIQUES, L. V PIRES, AND S. M. F. COZZOLINO, Iodo, in **Biodisponibilidade de nutrientes**, 4th ed., S. M. F. Cozzolino, Ed. Barueri: Manole LTDA, 2012, p. 767–795.
- [4] RUPÉREZ, P. Mineral content of edible marine seaweeds. **Food Chem**, v. 79, n. 1, p. 23–26, 2002.
- [5] COFRADES, S.; LÓPEZ-LOPEZ, I.; BRAVO, L.; RUIZ-CAPILLAS, C.; BASTIDA, S.; LARREA, M. T.; JIMÉNEZ-COLMENERO, F. Nutritional and antioxidant properties of different brown and red Spanish edible seaweeds. **Food Sci Technol Int**, v. 16, n. 5, p. 361–370, 2010.
- [6] BOCANEGRA, A., NIETO, A., BLAS, B. E SÁNCHEZ-MUNIZ, FJ. Dietas contendo uma alta porcentagem de algas Nori ou Konbu são bem aceitas e utilizadas eficientemente por ratos em crescimento, mas induzem diferentes graus de alterações histológicas no fígado e intestino. **Food Chem Toxicol**, v 41, n. 11, 1473-1480, 2003.
- [7] DAWCZYNSKI, C.; SCHÄFER, U.; LEITERER, M.; JAHREIS, G. Rodenas De La Rocha. Nutritional and toxicological importance of macro, trace, and ultra-trace elements in algae food products, **J Agr Food Chem**, v. 55, n. 25, p. 10470–10475, 2007

- [8] BESADA, V.; ANDRADE, J. M.; SCHULTZE, F.; GONZÁLEZ, J. J. Heavy metals in edible seaweeds commercialised for human consumption. **J Marine Syst**, v. 75, n. 1–2, p. 305–313, 2009.
- [9] VIDOTTI, E. C.; ROLLEMBERG, M. D. C. E. Algas: Da economia nos ambientes aquáticos à bioremediação e à química analítica. **Quim Nova**, v. 27, n. 1, p. 139–145, 2004.
- [10] MARINHO-SORIANO, E.; MOREIRA, W. S. C.; CARNEIRO, M. A. A. Some aspects of the growth of *Gracilaria birdiae* (Gracilariales, Rhodophyta) in an estuary in northeast Brazil. **Aquac. Int**, v. 14, p. 327–336, 2006.
- [11] ALFASSI, Z.B. Epithermal neutron activation analysis. *J.Radioanal. Nucl. Chem.*, v.90, p:151-165, 1985.
- [12] MAIHARA, A. M., MOURA, P.L.C., FÁVARO, D. I. T., VASCONCELLOS, M.B. Assessment of iodine in Brazilian duplicate portion diets and in table salt. **J.Radioanal. Nucl. Chem.** v. 278, p. 391-392, 2008.
- [13] KONIECZKA, Piotr; NAMIESNIK, Jacek. Quality assurance and quality control in the analytical chemical laboratory: a practical approach. CRC Press, 2018.
- [14] NAGATAKI, S. The average of dietary iodine intake due to the ingestion of seaweeds is 1.2 mg/day in Japan. **Thyroid**, v. 18, n. 6, p. 667–668, 2008.
- [15] FUKUSHIMA, M.; CHATT, A. Estimation of total as well as bioaccessible levels and average daily dietary intake of iodine from Japanese edible seaweeds by epithermal neutron activation analysis. **J Radioanal Nucl Ch**, v. 294, n. 3, p. 471–478, 2012.
- [16] IOM INSTITUTE OF MEDICINE REFERENCE INTAKES (DRIs) Estimated average requirements: Food and Nutrition Board, Institute of Medicine, National Academies, 2006. Disponível
<http://www.nal.usda.gov/fnic/DRI/DRI_Tables/recommended_intakes_individual.pdf.
Aceddado em 22jul.2018.